



Mark Scheme (Results)

Summer 2024

Pearson Edexcel International GCSE
In Mathematics B (4MB1) Paper 01R

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

Summer 2024

Question Paper Log Number P73495A

Publications Code 4MB1_01R_2406_MS

All the material in this publication is copyright

© Pearson Education Ltd 2024

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.
Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)

Abbreviations

- cao – correct answer only
- cso – correct solution only
- ft – follow through
- isw – ignore subsequent working
- SC – special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent

- awrt – answer which rounds to
- eeoo – each error or omission
- cas – Correct answer scores full marks (unless from obvious incorrect working)
- wr – working required

No working

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

With working

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question: eg. uses 252 instead of 255; follow through their working and deduct 2A marks from any gained provided the work has not been simplified. (Do not deduct any M marks gained.)

If there is a choice of methods shown, then award the lowest mark, unless the subsequent working makes clear the method that has been used

Examiners should send any instance of a suspected misread to review (but see above for simple misreads).

Ignoring subsequent work

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

Parts of questions

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

Question		Working	Answer	Mark	Notes
1		$\frac{75}{3000}$ or $\frac{0.075}{3}$ or $\frac{7.5}{300}$ or $\frac{0.75}{30}$ oe		2	M1 a 'correct' but un-simplified fraction (condone decimals in the fraction) eg $\frac{75 \times 10^{-3}}{3}$ or an answer of 0.025 or 2.5% or 2.5×10^{-2} oe eg 25×10^{-3}
			$\frac{1}{40}$		A1cao Do not ISW ignore any units eg $\frac{1mm}{40mm}$
		<i>cas</i>			Total 2 marks

Question		Working	Answer	Mark	Notes
2	(a)		5.143	1	B1
	(b)		6280	1	B1 allow 628×10 or 628×10^1 or 6.28×10^3 or 6.280×10^3 or 62.8×10^2 62.80×10^2 Do not accept 6.2800×10^3
					Total 2 marks

Question		Working	Answer	Mark	Notes
3			$4t^5v^2$	2	B2 (B1 for an expression in which 2 of the terms have been simplified correctly. eg $4t^5$ or $4t^5v$)
		<i>cas</i>			Total 2 marks

Question		Working	Answer	Mark	Notes
4	(a)		275	1	B1
	(b)		eg 'it would mean $n^2 = -9$ which is not possible' or 'all numbers are greater than 100' oe	1	B1 a correct reason (which needs to refer to a value other than 37) eg <ul style="list-style-type: none"> • $n^2 = -9$ or $n = \sqrt{-9}$ or $7n^2 = -63$ and idea this is not possible (eg n^2 cannot be negative or n or $n^2 \geq 0$ or needs to be a whole number/positive • numbers cannot be less than 100 • numbers are > 100 or ≥ 100 • numbers are ≥ 107 or > 107 • the smallest value is 107 • the first term is 107 and terms increase /don't decrease
					Total 2 marks

Question		Working	Answer	Mark	Notes
5		6 180 000 000 oe eg 6.18×10^9 or 61.8×10^8 or 20 600 000 oe eg 20.6×10^6 or 2.06×10^n where $n \neq 7$ or $2 \times 10^7 + 0.6 \times 10^6$ or $2 \times 10^7 + 6 \times 10^5$ oe		2	M1 allow for a correct numerator in any form or for a correct answer in any form or for 2.06×10^n where n is an integer $\neq 7$ or for a correct sum with the fraction removed
			2.06×10^7		A1 Do not ISW mark their answer on the answer line or if no answer on the answer line their final answer.
		<i>cas</i>			Total 2 marks

Question	Working	Answer	Mark	Notes
6	$6a^4c$ or $46656a^{24}c^6$ or $36^3a^{24}c^6$ or $6^6a^{24}c^6$		2	M1 for correctly applying the power of 3 or the power of a 0.5 to all 3 terms or for an expression of 3 terms in the form ma^pc^q with 2 of m, p and q correct or $a^{12}c^3$
		$216a^{12}c^3$		A1 ISW Do not allow $6^3a^{12}c^3$
	<i>cas</i>			Total 2 marks

Question	Working	Answer	Mark	Notes
7	$\frac{9}{4}[\div]\frac{15}{7}$ oe		3	M1 for writing both mixed numbers as improper fractions. Do not need the division sign. Implied by the 2 nd M1
	eg $\frac{9}{4} \times \frac{7}{15}$ oe eg $\frac{135}{60} \times \frac{28}{60}$ oe			M1 for inverting the 2 nd fraction and showing multiplying or for writing the improper fractions over a common denominator
		$\frac{63}{60} = 1\frac{1}{20}$ or $\frac{21}{20} = 1\frac{1}{20}$		A1 dependent on both Method marks being awarded. For completion to the correct answer with full working shown. We need to see the improper fraction followed by the mixed number
	<i>wr</i>			Total 3 marks

Question	Working	Answer	Mark	Notes
8	$3 \times 12x^2$ or $36x^2$		3	M1 for differentiating the first term correctly
	$\left[\frac{16}{x^2} = \right] 16x^{-2}$			M1 for rewriting the second term as $16x^{-2}$ This can be seen anywhere including in an expression for dy/dx May be implied by $-2 \times 16x^{-3}$ oe
		$36x^2 - 32x^{-3}$		A1 oe eg $36x^2 - \frac{32}{x^3}$ Need not be simplified eg allow $3 \times 12x^2 - 2 \times 16 \times x^{-3}$
	<i>cas</i>			Total 3 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

9		39.5, 40.5, 2.15, 2.25, 0.55, 0.65		3	M1 one correct bound allow 40.4 $\dot{9}$ for 40.5, 2.24 $\dot{9}$ for 2.25, 0.64 $\dot{9}$ for 0.65
		$\frac{UB_a}{LB_c - UB_f}$ where $(40 < UB_a \leq 40.5)$ $(2.15 \leq LB_c < 2.2) - (0.6 < UB_f \leq 0.65)$			M1 where $40 < UB_a \leq 40.5$ and $2.15 \leq LB_c < 2.2$ and $0.6 < UB_f \leq 0.65$
			$\frac{40.5}{2.15 - 0.65} = 27$		A1 for 26.97 to 27 if all correct figures seen in the equation. Allow 40.4 $\dot{9}$ for 40.5 and 0.64 $\dot{9}$ for 0.65
		<i>wr</i>			Total 3 marks

Question	Working	Answer	Mark	Notes
10		$y \geq -1$	4	B1 oe accept > for \geq
		$x + 2y \leq 8$		B1 oe eg $y \leq 4 - \frac{x}{2}$ accept < for \leq
	eg $y = 2x + 3$ or $y - 2x = 3$ or $y - (1) = 2(x - (-1))$ oe			M1 for the correct equation for the line written in any form. eg with any sign ($= > < \geq \leq$)
		$y \leq 2x + 3$		A1 oe allow equivalent inequality eg $y - 2x \leq 3$ or $y - (1) \leq 2(x - (-1))$ accept < for \leq
		SC if $y \leq -1$ and $x + 2y \geq 8$ and $y \geq 2x + 3$ score B2. Allow correct sign without the =		
				Total 4 marks
	<i>cas</i>			

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

11	(a)		a circle centre A with radius 4 cm	1	B1 full circle. Radius 4cm (allow if closer to Y than to B)
	(b)		a correct bisector with arcs shown	2	<p>B2 for a correct bisector with 2 pairs of arcs Allow if it is only one side of the line AB (use overlay) (NB may use the circle as one of the pair of arcs) (B1 for a bisector without the arcs or only one pair drawn or correct arcs without bisector drawn (must cross with in the lines on overlay or would if they were extended))</p>
	(c)		correct region indicated	1	<p>B1 dep on at least B1 being awarded in (b) for the bisector drawn. For the correct region between the bisector and the circle identified. Allow if a partial circle around A is drawn which intersects with their bisector twice and at least B1 awarded for the bisector drawn. NB if they do not shade and just label R they need to put R both sides of the line AB (if they have drawn the line AB)</p>
Total 4 marks					

Q	Working	Answer	Mark	Notes
12	eg $x^2 + x^2 = 12.6^2$ or $[AC^2 =](0.5 \times 12.6)^2 + (0.5 \times 12.6)^2 [= 39.69 + 39.69^2 = 79.38]$ or eg $\sin 45 = \frac{AC}{12.6}$ or $\cos 45 = \frac{AC}{12.6}$ or $\cos 45 = \frac{6.3}{AC}$ or $\sin 45 = \frac{6.3}{AC}$ oe		4	M1 for a correct statement using Pythagoras or trigonometry to find side AB/AC NB do not award for $AB^2 + AC^2 = 12.6^2$ unless made clear $AB = AC$. Allow AB instead of AC May be Implied by the 2 nd M1
	$[x =] \sqrt{\frac{12.6^2}{2}} = \left[\sqrt{\frac{3969}{25}} = 8.909\dots \right]$ or $[AC =] \sqrt{(0.5 \times 12.6)^2 + (0.5 \times 12.6)^2} [= 8.9090\dots]$ or $[AB/AC =] 12.6 \times \sin 45$ or $12.6 \times \cos 45$ $[AC/AB =] \frac{6.3}{\cos 45}$ or $\frac{6.3}{\sin 45}$			M1 dependent on 1st M1 for a correct method to find AB or AC Allow for 8.9 or better (actual 8.909545...) or $\frac{63\sqrt{2}}{10}$ Allow 9 if $x^2 = 79.3\dots$ is seen
	$2 \times "8.909\dots" + 12.6$			M1 independent of the previous method marks. Allow $\frac{63 + 63\sqrt{2}}{5}$ For using $2 \times n + 12.6$ where $6.3 < n < 12.6$ If n is incorrect, working must be shown
		30.4		A1 awrt 30.4 (actual 30.41909...)
	<i>cas</i>			Total 4 marks

Qu	Working	Ans	Mar	Notes
----	---------	-----	-----	-------

13	$[QR =] \frac{8}{\tan 40} \text{ or } \tan 40 = \frac{8}{QR} \text{ or } \tan 50 = \frac{QR}{8} \text{ or } \frac{QR}{\sin 50} = \frac{8}{\sin 40} \text{ oe}$ $[PR =] \frac{8}{\sin 40} \text{ or } \frac{8}{\cos 50} (= 12.445\dots) \text{ or } \sin 40 = \frac{8}{PR} \text{ or } \cos 50 = \frac{8}{PR} \text{ oe}$		4 M1 for a correct method or expression to find QR or PR . Allow any letter(s) for QR/PR or mislabelling or equivalents eg $[QR =] 8 \tan 50 (= 9.534\dots)$ If no working shown allow for awrt 12 or awrt 10
	<p>eg $[QR =] \frac{8}{\tan 40} (= 9.534\dots)$ AND $[PR =] \frac{8}{\sin 40}$ or $\frac{QR}{\cos 40} (= 12.445\dots)$ oe or</p> <p>$[QR =] \frac{8}{\tan 40} (= 9.534\dots)$ AND $[PR =] \sqrt{8^2 + "9.534\dots"}^2 (= 12.445\dots)$ oe or</p> <p>$[PR =] \frac{8}{\sin 40} (= 12.445\dots)$ AND $[QR =] \sqrt{"12.445\dots"}^2 - 8^2 (= 9.534\dots)$ oe or</p> <p>area of $PTR = 0.5 \times \pi \times \left(\frac{"12.445\dots"}{2}\right)^2 [= 60.8\dots]$ or</p> <p>area of $PQR = \frac{1}{2} \times 8 \times "9.534\dots"$ or $\frac{1}{2} \times 8 \times "12.445\dots" \times \sin(90 - 40)$ or</p> <p>$\frac{1}{2} \times "12.445\dots" \times "9.534\dots" \sin 40 [= 38.1\dots]$</p>		<p>M1 dependent on first M1 being awarded for a correct method or expression to find QR AND PR or $0.5PR$. Allow equivalent expressions eg those allowed for the 1st M1</p> <p>NB $\frac{PR}{\sin 90} = \frac{QR}{\sin 50} = \frac{8}{\sin 40}$ oe gains M1M1</p> <p>or for a correct method to find the area of the semicircle PTR or a correct method to find area of PQR using their PR (from correct working) and $PQ = 8$</p> <p>Allow numbers written to 1 dp.</p> <p>When finding the areas "12.445" or "9.534" must come from correct working.</p>
	$0.5 \times "9.534\dots" \times 8 + 0.5 \times \pi \times \left(\frac{"12.445\dots"}{2}\right)^2 \text{ or}$ $0.5 \times 8 \times "12.445\dots" \sin(50) + 0.5 \times \pi \times \left(\frac{"12.445\dots"}{2}\right)^2 \text{ or}$ $0.5 \times "12.445\dots" \times "9.534\dots" \sin(40) + 0.5 \times \pi \times \left(\frac{"12.445\dots"}{2}\right)^2$		<p>M1 correct method to find the whole area. If working is shown ft their PR(diameter) and/or PQ if clearly labelled or marked on the diagram or comes from correct working. Allow $\frac{"12.445\dots"}{2}$ or "6.22..." for the radius.</p>
		99	A1 awrt 99 or awrt 98
cas			Total 4 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

14	<p>eg $6x + 18y = 42$ or $10x + 30y = 70$ $6x + 10y = 20$ $18x + 30y = 60$</p> <p>or</p> <p>eg $3\left(\frac{14-6y}{2}\right) + 5y = 10$ or $3x + 5\left(\frac{14-2x}{6}\right) = 10$</p>		4	<p>M1 For balancing the equations (only condone one arithmetic error in multiplication). and correct operation to eliminate selected variable applied to all terms in their 2 equations)</p> <p>or writing x or y in terms of the other variable and correctly substituting to gain an equation in one variable</p>
		$x = -1.25$ or $y = 2.75$		<p>A1 oe one correct value dep on M1 Allow fractions eg $-\frac{7}{4}$ and $\frac{11}{4}$</p>
	<p>eg $2 \times "-1.25" + 6y = 14$ oe or $3x + 5 \times "2.75" = 10$ oe</p>			<p>M1 (dep) correct method to find second variable – could start process again or use substitution. Dependent on previous M mark being awarded. If the value used is incorrect you may need to check their answer if full working not shown.</p>
		$x = -1.25$ and $y = 2.75$		<p>A1 oe for both correct values dep on at least one of the method marks being awarded. If switched on the answer line allow if seen correct in working</p>
	wr			Total 4 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

15		$c^2 = \frac{3y+5}{8-y}$ oe or $c\sqrt{8-y} = \sqrt{3y+5}$		4	M1 for either squaring both sides to remove the square root or removing the denominator of the expression.
		$8c^2 - c^2y = 3y+5$ or $8-y = \frac{3y}{c^2} + \frac{5}{c^2}$ or $8-y = \frac{3y+5}{c^2}$ oe			M1 dependent on 1 st M1 being awarded. For squaring both sides to remove the square root and removing the terms in y from the denominator of the expression and expanding to gain a correct equation. Implies the previous M1
		$8c^2 - 5 = 3y + c^2y$ oe or $-c^2y - 3y = -8c^2 + 5$ oe or $c^2y + 3y = 8c^2 - 5$ oe or $y(-c^2 - 3) = -8c^2 + 5$ oe or $-\frac{3y}{c^2} - y = \frac{5}{c^2} - 8$ oe			M1 for collecting the y terms on one side of the equation with the other terms on the other side. Allow one sign error If the 2 nd M1 has not been awarded then ft their equation providing the equation has 4 distinct terms with exactly 2 in terms of y. Allow one sign error
			$y = \frac{8c^2 - 5}{3 + c^2}$		A1 oe eg $y = \frac{5-8c^2}{-c^2-3}$ or $y = \frac{8-\frac{5}{c^2}}{\frac{3}{c^2}+1}$ oe (NB: if the final answer is missing $y = \dots$ but is otherwise correct, award full marks if $y =$ a correct expression has been seen in the working otherwise do not ISW)
		<i>cas</i>			Total 4 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

16	eg $8 \times 4.5 = 3CP$ or $\frac{8}{3} = \frac{CP}{4.5}$ or $\frac{8}{3} = \frac{x+(x+3)}{4.5}$ $\frac{8}{3} = \frac{r+(r-3)}{4.5}$ oe		4	M1 for forming a correct equation. Using the chord theorem.
	$[CP =] \frac{8 \times 4.5}{3} [= 12]$ or $r = \frac{8 \times 4.5 + 3}{3 \times 2} [= 7.5]$ or $x = \frac{8 \times 4.5 - 3}{3 \times 2} [= 4.5]$			M1 for a correct expression for CP May be seen on diagram
	[circumference =] $\pi \times ("12" + 3)$ oe $2 \times \pi \times "7.5"$ or $2\pi \times ("4.5" + 3)$			M1 ft their CP if it comes from correct working or is clearly labelled or on the diagram. Allow 3.14... or $\frac{22}{7}$ for π
		47.1		A1 awrt 47.1 or 15π
	cas			Total 4 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

17	(a)	$\left[\begin{matrix} \vec{OB} \\ \end{matrix} \right] = \begin{pmatrix} 3 \\ -2 \end{pmatrix} - \begin{pmatrix} 5 \\ -9 \end{pmatrix}$ or $\begin{pmatrix} 3 \\ -2 \end{pmatrix} + \begin{pmatrix} -5 \\ 9 \end{pmatrix}$ or		2	M1 for a correct method to find \vec{OB} or may be implied by one correct value ie $\begin{pmatrix} -2 \\ \dots \end{pmatrix}$ or $\begin{pmatrix} \dots \\ 7 \end{pmatrix}$ or for $\begin{pmatrix} 7 \\ -2 \end{pmatrix}$ Must be seen on (a)
			$\begin{pmatrix} -2 \\ 7 \end{pmatrix}$		A1
	(b)	$\sqrt{(-2)^2 + 7^2} [= 7.28\dots]$ oe		2	M1 allow use of 2 rather than -2 and -7 rather than 7. May be implied by a correct answer or awrt 7.28 Condone missing brackets around negative values. Only ft their values from (a) or \vec{OB} found in (b) if working shown (allow \pm their values)
			$\sqrt{53}$		A1 ft their answer (as a simplified surd) to (a) or \vec{OB} found in (b) Do not ISW. This mark implies the M1 If no working is shown you may need to check.
		<i>cas</i>			Total 4 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

18	(a)		$30 < h \leq 40$	1	B1 condone \leq for $<$ and vice versa
	(b)	$5 \times 2 + 15 \times 26 + 25 \times 10 + 35 \times 24 + 45 \times 18$ (= 2300) $10 + 390 + 250 + 840 + 810$ (= 2300)		4	M2 for at least 3 correct products using midpoints with intention to add. (M1 for at least 3 products using frequency and a value within the interval with the intention to add. (allow use of upper/lower class bound) or for at least 3 correct products using midpoints without adding)
		$\frac{"10 + 390 + 250 + 840 + 810"}{80} \left[= \frac{"2300"}{80} \right]$			M1 dep on at least M1 being awarded
			28.75		A1oe allow $\frac{115}{4}$ or 28.7 or 28.8 or (29 from correct working)
		<i>cas</i>			Total 5 marks

Question	Working	Answer	Mark	Notes
19	$2x+3+5x-12+3x+1=32$ oe eg $10x-8=32$		5	M1 for setting up an equation. Implied by $x = 4$
	$x = \frac{32+8}{10} [= 4]$ oe or 7T vanilla = 13 oe			M1 rearranging to find a correct expression for x or for 7T vanilla
	$\frac{80}{360} \times 45$ (=10) oe			M1 Correct method to find the number for vanilla for 8Y eg $45 - \frac{280}{360} \times 45$
	$3 \times "4" + 1 - "10"$ or			M1 dependent on the 1 st and 3 rd M1 being awarded. Allow an answer of 3 or a correct method to find their difference. Ft their value of x and 10 if clearly labelled or come from correct working.
		3		A1 dependent on 1st and 3rd method mark being awarded. NB the number 3 can be gained from incorrect working, so need to check it. If no working is shown then award 5/5
				Total 5 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

20	(a)	$7+3y=2(4y-7-5)$ oe or $7+3y+2\times 5=2(4y-7)$ oe or $\frac{7}{2}+\frac{3}{2}y+5=4y-7$ oe		3	M1 for dealing with the fraction (of 0.5) correctly. Condone missing brackets if they have recovered.
		$3y-8y=-14-10-7$ oe eg $5y=31$ or $\frac{7}{2}+5+7=4y-\frac{3}{2}y$ oe			M1ft previous stage as long as at least 4 terms to deal with – for collecting terms in y on one side and number terms the other side
		<i>wr</i>	6.2		A1 dep on both the previous 2 Marks being awarded oe eg $\frac{31}{5}$
	(b)	$-35=8x-2\times -4.5$ oe or $x=\frac{A+2w}{8}$		2	M1 for a correct substitution into the given formula or for a correct rearrangement of the formula to make x the subject
		<i>cas</i>	-5.5		A1 oe eg $\frac{-44}{8}$ or $\frac{-11}{2}$
					Total 5 marks

Que	Working	Ans	M	Notes
-----	---------	-----	---	-------

21	<p>eg $\{2\pi r^2 \text{ and } 2\pi(3r)^2\}$ or $\left\{2\pi\left(\frac{R}{3}\right)^2 \text{ and } 2\pi R^2\right\}$ or</p> <p>$2R^2 + R^2 + 2r^2 - r^2$ or $2\pi R^2 + \pi R^2 + 2\pi r^2 - \pi r^2$ or</p> <p>$2(3)^2 + (3)^2 + 2(1)^2 - (1)^2$ or</p> <p>$2\pi(3)^2 + \pi(3)^2 + 2\pi(1)^2 - \pi(1)^2$ oe</p>		5	<p>M1 surface areas of both hemispheres seen (need not be added and may be part of an equation)</p> <p>Or for a correct formula for the total surface area. Allow $\{2\pi(nr)^2 \text{ and } 2\pi(mr)^2\}$ or $2(m)^2 + m^2 + 2(n)^2 - (n)^2$ or $2\pi(m)^2 + \pi m^2 + 2\pi(n)^2 - \pi(n)^2$ where $m = 3n$</p> <p>Allow use of other letters</p> <p>May be implied by the 2nd M1</p>
	<p>$2\pi r^2 + 2\pi(3r)^2 + \pi(3r)^2 - \pi r^2 = 567\pi$ oe or</p> <p>$2\pi R^2 + 2\pi\left(\frac{R}{3}\right)^2 + \pi R^2 - \pi\left(\frac{R}{3}\right)^2 = 567\pi$ oe or</p> <p>$2(3)^2 + (3)^2 + 2(1)^2 - (1)^2 : 567$</p> <p>$2\pi(3)^2 + \pi(3)^2 + 2\pi(1)^2 - \pi(1)^2 : 567\pi$</p>			<p>M1 a correct equation for the surface area of the solid</p> <p>eg $2(nr)^2 + 2(3nr)^2 + (3nr)^2 - (nr)^2 = 567$</p> <p>or $28\pi r^2 = 567\pi$ or $28r^2 = 567$</p> <p>or $2R^2 + 2\left(\frac{R}{3}\right)^2 + R^2 - \left(\frac{R}{3}\right)^2 = 567$</p> <p>Allow $2x^2 + 2(y)^2 + (y)^2 - x^2 = 567$</p> <p>$2(y)^2 + (y)^2 + 2(x)^2 - (x)^2 : 567$ where $y = 3x$</p> <p>This implies the 1st M1</p>
	[r =] 4.5 or [R =] 13.5			M1 for a correct value of the radius for either hemisphere. This implies the 2 nd M1
	$\frac{1}{2} \times \frac{4}{3} \pi \times "4.5"{}^3 + \frac{1}{2} \times \frac{4}{3} \pi \times "13.5"{}^3 \left[= \frac{243}{4} \pi + \frac{6561}{4} \pi \right]$			<p>M1 a fully correct method to find the volume of the solid dep on at least one of the previous Method marks being awarded and is equivalent to</p> <p>$\frac{1}{2} \times \frac{4}{3} \pi \times "m"{}^3 + \frac{1}{2} \times \frac{4}{3} \pi \times "3m"{}^3$ where m is a number</p> <p>(award 4 marks for 1701π)</p>
	wr	5340		A1 awrt to 5340 dependent on at least the 3 rd M1 mark being awarded from correct working (5343.8....)
				Total 5 marks

23	(a)	$6 \times (-4)^3 + k \times (-4)^2 - 26 \times -4 - 24 = 0$ or $6 \times (-4)^3 + 19 \times (-4)^2 - 26 \times -4 - 24$		2	M1 correct substitution of $x = -4$ into equation. Must be equal to zero (or implied by later working) or if substituting $x = -4$ and $k = 19$ into the expression we do not need it = 0 for this mark. Condone missing brackets around the -4
			$16k = 304$ or $k = \frac{304}{16}$ and $k = 19$ or $-384 + 304 + 104 - 24 = 0$ and shown		A1 completion to show clearly that $k = 19$ either by completing equation correctly (must see at least one line of working after the first line) and $k = 19$ or for $-384 + 304 + 104 - 24 = 0$ and comment (eg shown, or #)
	(b)	$(6x^2 \dots\dots\dots)$		4	M1 for a start to find the quadratic factor.
		$(6x^2 - 5x - 6)$			M1 for a correct 3 term quadratic
		$(3x+2)(2x-3)$ or $(3x+2)(2x-3)(x+4)$			M1 dependent on the 2 nd M1 being awarded for correct factorisation of the quadratic. Do not allow fractions or decimals eg $(x-1.5)$ or $\left(x+\frac{2}{3}\right)$ (ie) or a correct use of the quadratic formula. Implied by $-\frac{2}{3}$ and $\frac{3}{2}$ as two of the solutions (Allow -0.67 or better for $-\frac{2}{3}$)
			$-4, -\frac{2}{3}, \frac{3}{2}$		A1 oe $(-0.67$ or better for $-\frac{2}{3})$ dep on all 3 method marks being awarded. Do not ISW. Mark the answer on the answer line. If no answer on the answer line mark the final line of their working.
		<i>wr</i>			Total 6 marks

Question	Working	Answer	Mark	Notes
----------	---------	--------	------	-------

24	eg $\frac{0.5N+12.5}{N}$ or $\frac{N+25}{2N}$ or $\frac{p}{2p-25}$ or $\frac{25+y}{2y+25}$ oe		6	M1 for the probability that the first sweet is pink (N = number of sweets, p = number of pink sweets, y = number of yellow sweets) It must be seen as an expression in one letter and may be see in an equation
	eg $\frac{0.5N+12.5}{N} \times \frac{0.5N+11.5}{N-1} = \frac{7}{19}$ or $\frac{N+25}{2N} \times \frac{N+23}{2N-2} = \frac{7}{19}$ or $\frac{p}{2p-25} \times \frac{p-1}{2p-26} = \frac{7}{19}$ or $\frac{y+25}{2y+25} \times \frac{y+24}{2y+24} = \frac{7}{19}$ oe			M1 for a correct equation for 2 pink sweets. Must be expressed using one variable. This mark implies the 1 st M1
	eg $2.25N^2 - 235N - 2731.25 = 0$ or $9N^2 - 940N - 10925 = 0$ or $9p^2 - 695p + 4550 = 0$ or $9y^2 - 245y - 7200 = 0$			M1 a correct 3 term quadratic for N or p or y (allow any letter) condone missing =0
	eg $(9N + 95)(N - 115)$ or $(9p - 65)(p - 70)$ or $(9y + 160)(y - 45)$ oe			M1 dep on 2 nd M1 for a correct method to solve their 3 term quadratic – if factorising allow brackets that multiply giving 2 correct terms, if formula used then allow one error. Working must be shown if the equation is incorrect. Seeing or using $N = 115$ or $p = 70$ or $y = 45$ implies the 3 rd and 4 th Method marks
	$\frac{45}{115} \times \frac{44}{114}$			M1 for a correct calculation for the probability of 2 yellow sweets May be implied by a correct answer
	<i>wr</i>	$\frac{66}{437}$		A1 oe eg 0.151(029...) The correct answer will gain full marks if at least 2 method marks have been awarded.
	PTO for SC			

	SC for yellow is 25 more than pink M0M1M1M1M0A0			
	$\frac{0.5N - 12.5}{N} \times \frac{0.5N - 13.5}{N - 1} = \frac{7}{19} \text{ or}$ $\frac{N - 25}{2N} \times \frac{N - 27}{2N - 2} = \frac{7}{19} \text{ or}$ $\frac{y - 25}{2y - 25} \times \frac{y - 26}{2y - 26} = \frac{7}{19} \text{ or}$ $\frac{p}{2p + 25} \times \frac{p - 1}{2p + 24} = \frac{7}{19}$			M1 for a correct equation for 2 pink sweets. Must be expressed using one variable.
	$2.25N^2 + 240N - 3206.25 = 0 \text{ or}$ $9N^2 + 960N - 12825 = 0$ $9y^2 + 255y - 7800 = 0$ $9p^2 + 705p + 4200 = 0$			M1 a correct quadratic for N, p or y condone missing =0
	$[N =] \frac{240 \pm \sqrt{(240)^2 + 4 \times 2.25 \times 3206.25}}{2 \times 2.25} \text{ or}$ $[N =] \frac{-960 \pm \sqrt{960^2 + 4 \times 900 \times 12825}}{2 \times 9} \text{ or}$ $[y =] \frac{255 \pm \sqrt{255^2 + 4 \times 9 \times 7800}}{2 \times 9}$ $[p =] \frac{-705 \pm \sqrt{705^2 - 4 \times 9 \times 4200}}{2 \times 9} \text{ or}$			M1 dep on 2 nd M1 for a correct method to solve their 3 term quadratic – if factorising allow brackets that multiply giving 2 correct terms, if formula used then allow one error. Working must be shown
	wr			Total 6 marks

Quest	Working	Ans	Mark	Notes
25	$0.5 \times 8 \times 11 \times \sin 115 (= 39.877\dots)$		5	M1 for a correct method to find the area of triangle ABC
	$AC^2 = 11^2 + 8^2 - 2 \times 11 \times 8 \cos 115 (= 259.38\dots)$ or $AC = \sqrt{11^2 + 8^2 - 2 \times 11 \times 8 \cos 115} (= 16.105\dots)$			M1 for a correct method to find the length of AC^2 or AC
	$\angle CAD = \sin^{-1} \left(\frac{\sin(53) \times 15}{"16.105\dots"} \right) [= 48.059\dots]$ AND $\angle ACD = 180 - "48.059" - 53 (= 78.94\dots)$ or $16.105\dots^2 = AD^2 + 15^2 - 2 \times AD \times 15 \cos 53$ or $AD^2 = 16.105\dots^2 + 15^2 - 2 \times 16.105\dots \times 15 \cos "78.9\dots"$ $\left\{ \sqrt{"16.105\dots"}^2 - 11.979\dots^2 \text{ or } "16.105\dots" \cos \left[\sin^{-1} \left(\frac{\sin(53) \times 15}{"16.105\dots"} \right) \right] [= 10.76\dots] \right\}$ AND $\left\{ \sqrt{15^2 - 11.979\dots^2} \text{ or } 15 \cos 53 [= 9.027\dots] \right\}$			M1 for a correct method to find the angle ACD or Finding AD^2 NB $2 \times 15 \cos 53 = 18.054\dots$ the length AD (19.79...) For splitting triangle ACD into 2 triangles with perpendicular CE where E is on AD and finding the lengths AE (10.76...) and ED (9.027...)
	$\frac{1}{2} \times 15 \times "16.105\dots" \sin("78.94\dots") + "39.877\dots"$ or $\frac{1}{2} \times "19.79\dots" \times 15 \sin(53) + "39.877\dots"$ or $\frac{1}{2} \times "16.105\dots" \times "19.79\dots" \left(\frac{\sin(53) \times 15}{"16.105\dots"} \right) + "39.877\dots"$ or $\frac{1}{2} \times 15 \sin 53 \times "10.76\dots" + \frac{1}{2} \times 15 \sin 53 \times "9.027\dots" + "39.877\dots"$			M1 dependent on all 3 M marks being awarded. For a fully correct method to find the area NB $15 \sin 53 = 11.979\dots$
		158.42		A1 for awrt 158 (allow 159)
	<i>cas</i>			Total 5 marks

Question	Working	Answer	Mark	Notes
26	eg $\frac{6(x+3)+4(x-2)}{(x-2)(x+3)}$		5	M1 writing the addition part as a correct fraction over a common denominator – need not be expanded and may be 2 separate fractions. Allow one sign error in numerator if expanded
	eg $\frac{10(x+1)}{(x-2)(x+3)}$ or $\frac{10x+10}{x^2+x-6}$ oe			A1 a correct single fraction with numerator and denominator simplified – numerator and / or denominator may be factorised. This implies the 1st M1
	$\frac{(5x-5)(x-2)}{(x+1)(x-1)}$ or $\frac{(5x-10)(x-1)}{(x+1)(x-1)}$			M1 for numerator or denominator factorised correctly into 2 brackets
	$\frac{5(x-1)(x-2)}{(x+1)(x-1)}$ or $\frac{5(x-2)}{(x+1)}$ oe			M1 for numerator and denominator fully factorised correctly including factor of 5 taken out (could be implied by further cancelling) This implies the 2 nd M1
		$\frac{50}{x+3}$		A1 dep on M3
ALT	$\frac{(5x-5)(x-2)}{(x+1)(x-1)}$			M1 for numerator or denominator factorised correctly
	$\frac{5(x-1)(x-2)}{(x+1)(x-1)}$ or $\frac{5(x-2)}{(x+1)}$			M1 for numerator and denominator fully factorised correctly including factor of 5 taken out (could be implied by further cancelling)
	$\frac{30(x+3)+20(x-2)}{(x+1)(x+3)}$ oe			M1 for multiplying each part and writing the addition as a correct fraction over a common denominator – need not be expanded and may be 2 separate fractions. Allow one sign error in numerator if expanded
	$\frac{50x+50}{(x+1)(x+3)}$ or $\frac{50(x+1)}{(x+1)(x+3)}$			M1 for numerator and denominator fully factorised correctly (could be implied by further cancelling)
		$\frac{50}{x+3}$		A1 dep M3
	wr			Total 5 marks

