Please check the examination details belo	ow before entering your candidate information
Candidate surname	Other names
Pearson Edexcel Inter	
Friday 7 June 2024	
Morning (Time: 2 hours)	Paper reference 4PM1/02R
Further Pure Mat	hematics
Calculators may be used.	Total Marks

## **Instructions**

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Without sufficient working, correct answers may be awarded no marks.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- You must NOT write anything on the formulae page.
   Anything you write on the formulae page will gain NO credit.

## **Information**

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.

Turn over ▶





#### **International GCSE in Further Pure Mathematics Formulae sheet**

#### Mensuration

Surface area of sphere =  $4\pi r^2$ 

Curved surface area of cone =  $\pi r \times \text{slant height}$ 

Volume of sphere =  $\frac{4}{3}\pi r^3$ 

#### **Series**

## **Arithmetic series**

Sum to *n* terms,  $S_n = \frac{n}{2} [2a + (n-1)d]$ 

## **Geometric series**

Sum to *n* terms, 
$$S_n = \frac{a(1-r^n)}{(1-r)}$$

Sum to infinity,  $S_{\infty} = \frac{a}{1-r} |r| < 1$ 

#### **Binomial series**

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \dots + \frac{n(n-1)\dots(n-r+1)}{r!}x^r + \dots$$
 for  $|x| < 1, n \in \mathbb{Q}$ 

#### **Calculus**

## **Quotient rule (differentiation)**

$$\frac{\mathrm{d}}{\mathrm{d}x} \left( \frac{\mathrm{f}(x)}{\mathrm{g}(x)} \right) = \frac{\mathrm{f}'(x)\mathrm{g}(x) - \mathrm{f}(x)\mathrm{g}'(x)}{\left[\mathrm{g}(x)\right]^2}$$

## **Trigonometry**

### Cosine rule

In triangle ABC:  $a^2 = b^2 + c^2 - 2bc \cos A$ 

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B \qquad \sin(A-B) = \sin A \cos B - \cos A \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$cos(A + B) = cos A cos B - sin A sin B$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

### Logarithms

$$\log_a x = \frac{\log_b x}{\log_b a}$$



# Answer all ELEVEN questions.

# Write your answers in the spaces provided.

## You must write down all the stages in your working.

1	701	1 4	4.*
	Ina	allo arotic	equation
	1110	uuaurain	Cuualion

$$kx^2 - (2k+6)x + 16 = 0$$

has equal roots.

Find the two possible values of k

**(4)** 

	(Total for Question 1	is 4 marks)



2	(a)	Expand $\frac{2}{\sqrt{1+3x}}$ in ascending powers of x up to and including the term in $x^3$ Express each coefficient as a fraction in its simplest terms where appropriate.	(4)
	(b)	State the range of values of $x$ for which the expansion is valid.	(1)



 $\mathbf{3}$  O, A and B are fixed points such that

$$\begin{vmatrix} \overrightarrow{OA} \end{vmatrix} = 3\sqrt{5}$$
  $\overrightarrow{AB} = \mathbf{i} + 3a\mathbf{j}$   $\overrightarrow{OB} = 7\mathbf{i} + 2a\mathbf{j}$ 

Given that a > 0

(a) find the value of a

(4)

(b) Hence find a unit vector parallel to  $\overrightarrow{OA}$ 

**(2)** 




4	$f(x) = px^3 + qx^2 - 37x - 12q \text{ where } p \text{ and } q \text{ are constants.}$ When $f'(x)$ is divided by $(x + 2)$ the remainder is $-33$	
	Given that $(x + 5)$ is a factor of $f(x)$ (a) (i) show that $p = 2$ (ii) find the value of $q$	
	(b) Hence, use algebra to factorise $f(x)$ completely.	(6)
	(c) Hence solve the equation $f(x) = 0$	(2)



**(6)** 

5 The force F newtons between two magnetic poles is given by the formula

$$F = \frac{3}{20r^2}$$

where r is the distance, in centimetres, between the poles.

The distance between the two poles is increasing at a constant rate of 0.7 cm/s

Find the rate of change of F, in newtons/s to 3 significant figures, when the distance between the poles is  $2.8~\mathrm{cm}$ 






Figure 1

Figure 1 shows part of the graph of the curve with equation  $y = x + 2^{-(4x+1)}$ 

By drawing a suitable straight line on the graph, obtain an estimate, to one decimal place, of the roots of the equation  $\log_2(8-3x) + 4x = 0$  in the interval  $-2 \le x \le 6$ 

**(7)** 






Figure 2

Figure 2 shows a sketch of part of the curve C with equation

$$y = \frac{x^2 - 1}{4x + 5} \quad \text{where} \quad x \neq -\frac{5}{4}$$

(a) Write down the equation of the asymptote to C that is parallel to the y-axis.

(1)

The line *l* is the normal to *C* at the point where x = -1

(b) Find an equation of l

**(7)** 

The line l meets C again at the point D

(c) Find the coordinates of D

(6)



Question 7 continued	





8 The sum of the first 2 terms of a geometric series G is 360 The sum of the 2nd and 3rd terms of G is 288

The *n*th term of G is  $U_n$ 

(a) Show that  $U_n = A \left(\frac{4}{5}\right)^{n-1}$  where A is an integer to be found.

(7)

(b) Explain why G is convergent.

(1)

(c) Hence find the sum to infinity of G

(2)

(d) Find the least number of terms for which the sum is greater than 978

**(4)** 

| <br> |
|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|
| <br> | <br> | <br> | <br> | <br> | <br> |      |      | <br> |      | <br> |      |
|      |      |      |      | <br> | <br> |      | <br> |      | <br> |      |      |      |      |      |      | <br> | <br> | <br> |      |      |      |      |      |      |
| <br> |





Question 8 continued








Figure 3

Figure 3 shows a sketch of part of the curve S with equation  $y = -2e^{3x} + 4$  and the line L

The curve S has intersections with the line L at the points A and B with x coordinates x = -1 and x = 0 respectively.

The finite region bounded by S and L is shown shaded in Figure 3

Use calculus to find the exact area of this region.

Give your answer in the form  $\frac{a + be^{-c}}{c}$  where a, b and c are integers to be found.

(8)

2	7
_	4



Question 9 continued	








Figure 4

Figure 4 shows a right pyramid *ABCDV* 

The base of the pyramid is a rectangle where,

$$AB = DC = 2x$$
 cm  $AD = BC = x$  cm

The edges VA, VB, VC and VD are all of equal length.

The angle between VA and ABCD is 45°

(a) Show that 
$$VA = \frac{\sqrt{10}}{2} x \text{ cm}$$

(3)

(b) Find in cm, the exact height of the pyramid in terms of x

(2)

Find, in degrees to one decimal place,

(c) the size of angle VBA

**(2)** 

(d) the size of the obtuse angle between the plane AVC and the plane BVD

**(4)** 

Given that the volume of the pyramid is  $9\sqrt{5} \text{ cm}^3$ 

(e) find the value of x

(2)





	$\Diamond \Diamond$
	XX.
$\longleftrightarrow$	$\times\!\!\times$
$\longleftrightarrow$	$\times\!\!\times$
$\leftrightarrow\!$	$\times\!\!\times$
DO NOT WRITE	$\times\!\!\times$
$\sim$	$\times\!\!\times$
	$\times\!\!\times$
	88.
$\times \times$	88.
	88.
X	88.
$\times$	$\Diamond \Diamond$
$\times \omega$	$\otimes$
$\times\!\times\!\times$	$\Diamond \Diamond$
$\times \nabla$	$\Diamond \Diamond$
$\times\!\times\!\times$	$\Diamond \Diamond$
A 55	XX.
8.55	XX.
O THE	XX.
	XX.
	$\times$
	$\times\!\!\times$
	$\times\!\!\times$
$\bigcirc$ 777	$\times\!\!\times$
	$\times\!\!\times$
<b>○</b>	$\times\!\!\times$
0 <del>99</del>	$\times\!\!\times$
	$\times \times$
	$\times\!\!\times$
SIMTMIS	88.
$\times \times \times$	XX.
$\times \mathbb{Z}^{K}$	XX.
XXX	XX.
XXX	88.
$\times$	88
XXX	88
X	(X)
$\times$	ΦŎ.
Xida	$\Diamond \Diamond$
$\times$	$\Diamond \Diamond$
X	$\Diamond \Diamond$
AREA	$\Diamond \Diamond$
$\times$	$\Diamond \Diamond$
$\times$	$\Diamond \Diamond$
$\times\!\!\times\!\!\!\times$	$\Diamond \Diamond$
$\times\!\!\times\!\!\times$	$\Diamond \Diamond$
$\times\!\!\times\!\!\times$	$\Diamond \Diamond$
$\times\!\times\!\times$	$\Diamond \Diamond$
$\times\!\!\times\!\!\times$	$\propto \sim$
$\times\!\!\times\!\!\times$	$\times$
$\times\!\!\times\!\!\times$	$\times\!\!\times$
$\times\times\times$	XX.
$\sim\sim$	XX.
	$\times \times$
	$\otimes$
	$\otimes$
	$\otimes$
	$\overset{\circ}{\otimes}$
	×
Q	
DC.	
DO	
DOI	
0	
0	
0	
0	
0	
0	
0	
0	
NOT WRIT	
NOT WRIT	
NOT WRIT	
NOT WRITE	
NOT WRITE	
NOT WRITE IP	
NOT WRITE IN THIS	
NOT WRITE IN THIS A	
NOT WRITE IN THIS	

					é		
					è		
			ì				
			2	۹	7	ę	
þ	K	>	ζ	3	Ζ	3	
>	S	>	Ś		Ž	Š	
>	8	3	ξ			į	
>	S	3	Š	4			֡
>	3	3	S	d			֡
	5	3	S	d			
>	3	3	S	d			
	3	3	3		2	2	
>	3	3	3		2	2	
	3	3	3		2	2	
	3	3				2	
>		3				2	
>	\ \ \	?					
		3					
		?				2	
>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>>		3					
		\ \ \ \					
		3					
		?					
		3					

Question 10 continued	





11 (a) Using a formula on page 2 show that  $\cos 2A = 2\cos^2 A - 1$ 



(b) Hence show that  $(2\cos^2 A - 1)^2 = \frac{\cos 4A + 1}{2}$ 



The curve with equation  $y = \frac{\sin 2x}{2} + \frac{\left(2\cos^2 x - 1\right)^2}{2} + \frac{1}{8}$  has a stationary point P in the range  $0 \le x \le \frac{\pi}{6}$ 

(c) Find the exact coordinates of P








Question 11 continued	
	(Total for Question 11 is 12 marks)
	TOTAL FOR PAPER IS 100 MARKS

